

AVIFAUNA OF THE NORTHERN CORDILLERA CENTRAL OF THE ANDES, COLOMBIA

Andrés M. Cuervo¹, Paulo C. Pulgarín², Diego Calderón-F.², José M. Ochoa-Quintero³, Carlos A. Delgado-V.², Alejandro Palacio², Jorjany M. Botero², & Wilmar A. Múnera²

¹Department of Biological Sciences and Museum of Natural Science, Louisiana State University, 119 Foster Hall, Baton Rouge, LA 70803, U.S.A. E-mail: acuerv1@lsu.edu

²Instituto de Biología, Universidad de Antioquia, A. A. 1226, Medellín, Colombia.

³Institute for Biodiversity and Ecosystem Dynamics, Amsterdam University, Kruislaan 318, 1098 SM Amsterdam, The Netherlands.

Resumen. – **Avifauna del norte de la Cordillera Central de los Andes, Colombia.** – Entre 1999 y 2006, realizamos inventarios de aves mediante observaciones de campo, grabaciones sonoras y colecta de especímenes en el extremo norte de la Cordillera Central Andina, departamento de Antioquia, Colombia. Registramos un total de 335 especies en 18 bosques de niebla localizados entre 1300 y 1915 m s.n.m. Entre las diez localidades más visitadas, Bodega Vieja (231 especies), Alto Chaquiral [= Arriero Antioqueño (202)], La Forzosa (199) y Santa Gertrudis (162) tuvieron la mayor riqueza. Presentamos una evaluación cualitativa de la abundancia relativa y uso de hábitat para cada especie. El ensamblaje de esta avifauna exhibe una mezcla de afinidades biogeográficas con elementos de los Andes del Norte, las tierras bajas y piedemonte del Chocó, Nechí-Magdalena, y Centro América. Además de esto, la cantidad de registros novedosos que encontramos contradicen la visión de la Cordillera Central como un área bien explorada en Colombia. Diecisésis especies prioritarias para la conservación tienen poblaciones importantes en estos bosques nublados; por ejemplo, acá reportamos las 16 localidades conocidas para el Arriero Antioqueño (*Lipaugus weberi*), cuyo rango de distribución está totalmente confinado a nuestra área de estudio. Las características únicas de esta avifauna resaltan la necesidad de una estrategia de conservación integral desde las comunidades locales para contrarrestar el impacto del cambio del paisaje que resulta de la expansión actual de la ganadería, caficultura, y cultivos ilícitos en la zona.

Abstract. – From 1999 to 2006, we conducted bird inventories based on observations, tape-recordings, and general specimen collecting in the northern end of the Andean Cordillera Central, Department of Antioquia, Colombia. We recorded 335 species in 18 cloud forests located between 1300 and 1915 m a.s.l. Among the ten localities more frequently visited, Bodega Vieja (231 species), Alto Chaquiral [= Arriero Antioqueño (201)], La Forzosa (199) and Santa Gertrudis (162) had the higher species richness. We present a qualitative assessment of relative abundance and habitat preferences for each species. The avifauna assemblage of this region exhibits mixed biogeographic affinities, with elements from the northern Andes, the lowlands and foothills of Chocó, Nechí-Magdalena, and Central America. Besides this, the outstanding new bird records we found contradict the view of the Cordillera Central as a relatively well explored area in Colombia. Sixteen species of conservation concern have healthy populations in these cloud forests; for instance, we report all the 16 localities known for the Chestnut-capped Piha (*Lipaugus weberi*) whose range is entirely confined to our study region. The unique features of this avifauna highlight the necessity of an integral, community-based conservation strategy to ameliorate the effect of landscape

changes resulting from the ongoing expansion of cattle ranching, coffee plantations, and illicit crops.
Accepted 7 June 2008.

Key words: Antioquia, Chocó, Nechí lowlands, Porce river, northwestern South America, premontane forests.

INTRODUCTION

The cloud forests of the Andes in northwestern South America are recognized for their extremely high levels of avian diversity and endemism (Stotz *et al.* 1996, Rahbek & Graves 2001). The peaks of this pattern lie along the eastern slope of the Andes from southern Peru to southern Colombia (Rahbek & Graves 2001, Hawkins *et al.* 2007). However, many more localities in that stretch of the Andes are better known, particularly in eastern Ecuador and Peru (Terborgh 1975, Stotz *et al.* 1996, Pitman *et al.* 2002), than other portions with continuous cloud forests along altitudinal gradients such as the unexplored areas of the Andes of Colombia (van Velzen 1992).

The northern Andes attain their higher complexity in Colombia where the main range split in three cordilleras: Occidental, Central and Oriental (Fig. 1). Each cordillera has a unique geological history and distinctive topographic, climatic and ecological features (McCourt *et al.* 1984, van Velzen 1992, Gregory-Wodzicki 2000, IGAC 2003); for instance, the Cordillera Central (Fig. 1) is the oldest (c. 65,000,000 years) and the highest on average, of the three (McCourt *et al.* 1984, Gregory-Wodzicki 2000). At its northern end, the Cordillera Central smoothly terminates in the very wet slopes and foothills of the Nechí and Porce valleys, and the Remedios-Segovia low pass to Serranía de San Lucas (Fig. 1). Ornithological studies conducted since 1999 have suggested that the avifauna of the premontane forests of the northern Cordillera Central is one of the most intriguing of northern South America. First, taxa from multiple

biogeographic areas converge to this region (Cuervo *et al.* in press). Second, several new species-level taxa, like the Chestnut-capped Piha (*Lipaugus weberi*), have been found in such a small area (Cuervo *et al.* 2001). Lastly, populations of threatened, poorly known, and rare cloud forest birds have been found to be relatively common and to thrive in its fragmented landscape (Cuervo & Restrepo 2007).

Here we report a comprehensive list of the avifauna of the cloud forests of the northern Cordillera Central as a result of our studies in 18 localities of the region from 1999 to 2006. Our goal is to provide baseline information for ongoing and future biodiversity research and conservation action in the region.

STUDY AREA AND METHODS

Our study area encompasses the northernmost end of the Cordillera Central of the Colombian Andes, consisting of 18 premontane wet forests (Espinol 1992) located between 1300–1915 m in the upper Nechí and lower Porce valley, municipalities of Anorí and Amalfi, Department of Antioquia, Colombia (Fig. 1, Table 1). The landscape consists of a collage of multiple fragments of mature cloud forest, including several large tracts of continuous forest, embedded in a matrix of pastures and, to a lesser extent, coffee plantations (Cuervo & Restrepo 2007).

Since 1999 we have been conducting opportunistic observations, recordings of bird vocalizations, and general specimen collecting that complemented a more comprehensive systematic survey focused on the nine localities reported by Cuervo & Restrepo (2007),




FIG 1. Map of northwestern South America depicting terrain above 1000 m elevation in gray. The location of our study area in the northern Cordillera Central is delimited by a square and shown in detail in the right panel. Open circles represent the location of study sites and numbers within correspond to the locality list in Table 1.

plus La Forzosa (Table 1). The systematic samplings consisted in 17–23 days of mist-netting in the interior and forest edge at each of this subset of nine localities. Copies of all recordings were archived at Banco de Sonidos Animales, Instituto Alexander von Humboldt, and many have been published in a sound guide (Álvarez *et al.* 2007). Voucher specimens were taken whenever possible and deposited in the Instituto de Ciencias Naturales (ICN), Museo Universidad de Antioquia (MUA), and Instituto Alexander von Humboldt (IAvH). Details on habitats, vegetation, sampling strategies, dates, and co-workers have been

reported elsewhere (Cuervo *et al.* 2005, Cuervo & Restrepo 2007).

RESULTS AND DISCUSSION

We recorded a total of 335 species in the cloud forests of the northern tip of the Cordillera Central (Appendix 1), a relatively high number for a local montane avifauna in such narrow elevation range. For example, Renjifo (1999) reported a comprehensive list of 243 species in a fragmented landscape of similar area in the western slope of this same cordillera. The high species richness of the north-

TABLE 1. Geographic coordinates (north latitude, west longitude), elevation range (m a.s.l.), accumulated number of days, and number of species recorded in 18 localities of the northern Cordillera Central. Locality numbers as mapped in Figure 1.

Localities	Coordinates	Elevation (m)	Fieldwork days	Species richness
1. Finca Los Canales*	6°49',75°05'	1810–1880	22	135
2. La Secreta*	6°49',75°06'	1820–1915	20	157
3. Bosque Guayabito*	6°52',75°06'	1700–1815	36	179
4. Bosque Las Ánimas*	6°56',75°01'	1500–1550	19	148
5. Bosque de El Abuelo-Escuela Las Ánimas*	6°56',75°00'	1505–1610	19	164
6. Bosques de Santa Catalina	6°57',75°02'	1500–1540	5	106
7. Bodega Vieja*	6°58',75°03'	1300–1550	34	231
8. Alto El Chaquiral* [=Arrierito Antioqueño]	6°58',75°08'	1650–1735	22	202
9. Reserva La Forzosa	6°59',75°08'	1550–1740	36	198
10. Finca Macondo	7°00',75°05'	1400–1600	5	73
11. Filo Largo	7°03',75°10'	1475–1520	1	47
12. Mangas de Rosita	7°04',75°08'	1520–1550	1	34
13. Mampuestos*	7°04',75°10'	1445–1495	17	136
14. Quebrada La Nutria	7°04',75°11'	1425–1440	1	39
15. La Condena	7°06',75°06'	1600–1750	3	59
16. Alto La Serrana	7°06',75°08'	1630–1700	5	84
17. Camino Anorí-Santa Gertrudis	7°06',75°09'	1600–1700	4	83
18. Santa Gertrudis*	7°08',75°09'	1420–1470	23	162

*One of the nine localities reported in Cuervo & Restrepo (2007), that along with La Forzosa (9) were more thoroughly sampled.

ern slope of the Cordillera Central is consistent with a peak in avian diversity expected at middle elevations in the tropical Andes, due to an area effect and influx of species from the lowlands (Rahbek 1997, Kattan & Franco 2004); this pattern, however, can only be tested when comparable data from the adjacent lowlands and highlands become available. Of the ten localities more thoroughly sampled, species richness was higher in Bodega Vieja, Bosque Guayabito, Alto Chaquiral [=Arrierito Antioqueño], La Forzosa, and Santa Gertrudis (Table 1). These are, overall, medium to large forests (from 110 to more than 1000 ha) with lower disturbance levels than the other sites. In addition to cloud forest birds, we also recorded lowland and open country species in the pastures,

young second growth, edges or gaps surrounding these forests (Appendix 1). We predict that, when properly sampled, La Condena and Macondo, two of the sporadically visited sites, will exhibit similar composition and richness values as the ones mentioned above given the large extension of forest they encompass.

The birds that assemble the avifauna of the northern Cordillera Central of Colombia show diverse biogeographic patterns (Cuervo et al. in press). To this region coincide the geographic ranges of taxa, at the species and subspecies level, characteristic of the northern Andes (e.g., *Phaethornis guy emiliae*, *Rhynchocyclus fulvipectus*, *Pyroderus scutatus occidentalis*, *Cypharhis nigrirostris*), the Nechí-Magdalena lowlands and foothills (e.g., *Capito hypoleucus*, *Hyloctistes*

subulatus cordobae, *Piprites chloris antioquia*e, *Habia gutturalis*), Chocó (e.g., *Androdon aequatorialis*, *Trogon comptus*, *Myiophobus phoenicomitra litiae*, *Diglossa indigotica*), or lowlands and foothills of Central America (e.g., *Philydor fuscipenne*, *Automolus rubiginosus saturatus*). Except for the adjacent northern Cordillera Occidental and Serranía de San Lucas (Fig. 1), no other avifauna in the northern Andes is expected to consist of such mixture of montane and trans-Andean lowland rainforest elements.

Overall, the species pool of the Cordillera Central of the Colombian Andes has been previously considered to be relatively well known. However, not only new species have been recently described (Cuervo *et al.* 2001, 2005; Donegan 2007), but also important range extensions have been reported (Beltrán & Kattan 2001). For example, we presented evidence for 36 unexpected species in the region that reinforced the pattern of mixture of biogeographical realms in this corner of South America (Cuervo *et al.* in press). That the avifauna of the Cordillera Central was presumed to be generally well known was based on the assumption that the various expeditions and recent works conducted in the coffee growing region in the departments of Antioquia, Caldas, Quindío, and Risaralda were representative of the whole range. Indeed, there are vast areas where little or no exploration has been conducted yet, such as the páramos and upper montane forests in northern Antioquia, the hill slopes of the dry Cauca canyon, the eastern slope from Antioquia to Tolima, and the southern portion from Parque Nacional Las Hermosas south to Parque Nacional Puracé. The finding of a distinct avifauna in its northern end implies that collecting surveys in the poorly-explored areas of this cordillera should be a priority for field ornithologists in Colombia.

Six IUCN threatened and ten near-threatened species have populations in the northern Cordillera Central (Appendix 1); many of

which were not even known to occur in this area at all such as *Tinamus osgoodi*, *Bangsia melanochlamys*, and *Chlorocrysa nitidissima*. Furthermore, *Lipaugus weberi* is exclusively known from the sixteen localities reported in this paper, twelve more since its description in 2001. Although this cotinga may indeed be restricted to the cloud forests of Anorí and Amalfi, ornithological exploration is poor in adjacent areas like Yolombó, Campamento, Yarumal, the northern Cordillera Occidental, and Serranía de San Lucas. Except for *Odontophorus hyperythrus*, *Pyrrilia pyrilia*, and *H. gutturalis*, the rest of species of conservation concern mostly range between 1000 and 2000 m, the altitudinal belt at which most forest cover have disappeared in Colombia (Etter & van Wyngaarden 2000). That many premontane birds locally extirpated in other portions of the Colombian Andes (see Renjifo 1999) have relatively large populations in our region is indicative of the mild history of disturbance of the northern end the cordillera. Instead of the extensive deforestation of premontane wet forests due to the expansion of coffee cultivations and cattle ranching that characterized the Colombian Andes in the 20th century, this region remained mostly forested in many parts due to a low local human population density and a gold-mining based economy. However, land use is rapidly changing due to the exhaustion of gold mines, the expansion of illicit crops and cattle ranching all over Colombia, and stable international coffee prices.

Bodega Vieja in Amalfi and Santa Gertrudis in Anorí are two key large forest blocks that still remain unprotected. By being located in the middle of a forest continuum from 1800 m a.s.l. to 150 m, Santa Gertrudis represents a unique opportunity for expanding the Bajo Cauca-Nechí reserve (45,000 ha) which currently protects the lower foothills and lowlands (150–850 m). There are no national parks in the northern Cordillera Central, but

three small reserves were recently established based on our studies: Bosque Guayabito (c. 1000 ha), La Forzosa (c. 350 ha), and Arrierito Antioqueño (i.e., Alto El Chaquiral, c. 110 ha). Although only the Anorí region was identified as an Important Bird Area (Franco & Bravo 2005), realistic conservation impact of the IBA program on the cloud forest habitats of the region is still to be seen. We believe that a network of reserves owned by local farmers would cradle long-term, sustainable conservation of the lower Porce and upper Nechi forests, which are now threatened by ongoing land use changes and increasing human colonization.

ACKNOWLEDGMENTS

We appreciate the hospitality of the local people of Anorí and Amalfi. Funding and equipment was provided by CORANTIOQUIA, the NSF-CREST program at University of Puerto Rico, Sigma Xi, Instituto Humboldt, Organization for Tropical Studies, Neotropical Bird Club, Idea Wild, Optics for the Tropics, the “Incentivos a la Investigación” program of Museo Universitario at Universidad de Antioquia, Cornell Laboratory of Ornithology, Asociación Calidris, and Barrick Museum of Natural History. CORANTIOQUIA granted research permits. We thank J. Toro for encouraging us to study the birds of this region. For their hard work and company in the field we are very grateful to many friends and students in Colombia acknowledged elsewhere. M. A. Echeverry-Galvis and C. D. Cadena made useful comments that improved the quality of the manuscript.

REFERENCES

- Álvarez, M., V. Caro, O. Laverde, & A. M. Cuervo. 2007. Guía sonora de las aves de los Andes colombianos/A guide to the birds sounds of the Colombian Andes. Banco de Sonidos Animales, Instituto Alexander von Humboldt & Cornell Laboratory of Ornithology, Ithaca, New York.
- Beltrán, W., & G. Kattan. 2001. First record of the Slaty-backed Nightingale-Thrush in the central Andes of Colombia, with notes on its ecology and geographical variation. Wilson Bull. 113: 134–139.
- Cuervo, A. M., & C. Restrepo. 2007. Assemblage and population-level consequences of forest fragmentation on bilateral asymmetry in tropical montane birds. Biol. J. Linn. Soc. 92: 119–133.
- Cuervo, A. M., P. Salaman, T. Donegan, & J. M. Ochoa. 2001. A new species of piha (Cotingidae: *Lipaugus*) from the Cordillera Central of Colombia. Ibis 143: 353–368.
- Cuervo, A. M., C. D. Cadena, N. Krabbe, & L. M. Renjifo. 2005. *Scytalopus stilesi*, a new species of tapaculo (Rhinocryptidae) from the Cordillera Central of Colombia. Auk 122: 445–463.
- Cuervo, A. M., P. C. Pulgarín, & D. Calderón-F. in press. New distributional bird data from the Cordillera Central of the Colombian Andes, with implications for the biogeography of northwestern South America. Condor.
- Donegan, T. 2007. A new species of brush finch (Emberizidae: *Atlapetes*) from the northern Central Andes of Colombia. Bull. Br. Ornithol. Club 127: 255–268.
- Espinal, L. S. 1992. Geografía ecológica de Antioquia: Zonas de vida. Editorial Léalon, Medellín, Colombia.
- Etter, A., & W. van Wyngaarden. 2000. Patterns of landscape transformation in Colombia, with emphasis in the Andean region. Ambio 29: 432–439.
- Franco, A. M., & G. A. Bravo. 2005. Áreas importantes para la conservación de las aves en Colombia. Pp. 117–282 in Áreas importantes para la conservación de las aves en los Andes tropicales: sitios prioritarios para la conservación de la biodiversidad. BirdLife International & Conservation International, Quito, Ecuador.
- Gregory-Wodzicki, K. M. 2000. Uplift history of the central and northern Andes: a review. Geol. Soc. Am. Bull. 112: 1091–1105.
- Hawkins, B. A., J. A. F. Diniz-Filho, C. A. Jaramillo,

- & S. A. Soeller. 2007. Climate, niche conservatism, and the global bird diversity gradient. *Am. Nat.* 170: S16–S27.
- IGAC. 2003. Atlas de Colombia. Instituto Geográfico Agustín Codazzi & Imprenta Nacional de Colombia, Bogotá, Colombia.
- Kattan, G., & P. Franco. 2004. Bird diversity along elevational gradients in the Andes of Colombia: area and mass effects. *Glob. Ecol. Biogeogr.* 13: 451–458.
- McCourt, W. J., J. A. Aspden, & M. Brook. 1984. New geological and geochronological data from the Colombian Andes: continental growth by multiple accretion. *J. Geol. Soc.* 141: 831–845.
- Pitman, N., D. K. Moskovits, W. S. Alverson, & R. Borman. 2002. Ecuador: Serranías Cofán-Bermejo, Sinangoe. Rapid Biological Inventories Report 3, The Field Museum, Chicago, Illinois.
- Rahbek, C. 1997. The relationship among area, elevation, and regional species richness in Neotropical birds. *Am. Nat.* 149: 875–902.
- Rahbek, C., & G. R. Graves. 2001. Multiscale assessment of patterns of avian species richness. *Proc. Natl. Acad. Sci.* 98: 4534–4539.
- Remsen, J. V., Jr., C. D. Cadena, A. Jaramillo, M. Nores, J. F. Pacheco, M. B. Robbins, T. S. Schulenberg, F. G. Stiles, & D. F. Stotz. 2008. A classification of the bird species of South America. American Ornithologists Union. Available on <http://www.museum.lsu.edu/~Remsen/SAC-CBaseline.html> [Version 24 April 2008].
- Renjifo, L. M. 1999. Composition changes in a sub-andean avifauna after long-term forest fragmentation. *Conserv. Biol.* 13: 1124–1139.
- Stotz, D. F., J. W. Fitzpatrick, T. A. Parker, III, & D. Moskovits. 1996. Neotropical birds: Ecology and conservation. Chicago Univ. Press, Chicago, Illinois.
- Terborgh, J. 1975. Bird species diversity on an Andean elevational gradient. *Ecology* 58: 1007–1019.
- van Velzen, H. P. 1992. Priorities for conservation of the biodiversity in the Colombian Andes. *Noved. Colomb.* 4 (Esp.): 1–32.

APPENDIX 1. Avifauna of the northern Cordillera Central of the Colombian Andes, indicating relative abundance, major habitats where the species were recorded, and type of evidence gathered per species. See Table 1 for codes and geographic data of the 18 localities studied. Taxonomy follows the A.O.U. South American Checklist Committee recommendations (Remsen *et al.* 2008).

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Tinamidae					
Black Tinamou (VU)	<i>Tinamus osgoodi</i>	9	SC	I	S,H
Little Tinamou	<i>Crypturellus soui</i>	7–10,18	U	III	R,H
Cracidae					
Sickle-winged Guan	<i>Chamaepetes goudotii</i>	1–5,7–9,13	FC	I,II,III	R,S,H
Crested Guan	<i>Penelope purpurascens</i>	7	SC	I	R,S,H
Wattled Guan (NT)	<i>Aburria aburri</i>	3,5,7–9,15–18	U	I	R,H
Speckled Chachalaca	<i>Ortalis guttata columbiana</i>	1–9,11,13,15–17	C	II,III	R,S,H
Odontophoridae					
Rufous-fronted Wood-Quail	<i>Odontophorus erythrops</i>	7,8	U	I	R,S,H
Chestnut Wood-Quail (NT)	<i>Odontophorus hyperythrus</i>	2,3,5,8,9,13,16,18	FC	I	R,S,H
Ardeidae					
Striated Heron	<i>Butorides striata</i>	7	X	IV	S
Cattle Egret	<i>Bubulcus ibis</i>	7–9	U	III	S
Cathartidae					
Turkey Vulture	<i>Cathartes aura</i>	1–9–12,13,18	C	V	S
Black Vulture	<i>Coragyps atratus</i>	1–18	C	V	S
King Vulture	<i>Sarcogyps papa</i>	2,6,7–10,18	U	V	S
Accipitridae					
Gray-headed Kite	<i>Leptodon cayanensis</i>	1	X	II	S
Swallow-tailed Kite	<i>Elanoides forficatus</i>	1–10,13,17,18	U	II,III,V	R,P,S,H
White-tailed Kite	<i>Elanus leucurus</i>	7,9	U	III	S
Semicollared Hawk (NT)	<i>Accipiter collaris</i>	7,8	SC	I,II	M,P,S,H
Sharp-shinned Hawk	<i>Accipiter striatus ventralis</i>	3,8,13	U	I,II	M,P
White Hawk	<i>Leucopternis albicollis</i>	8,9	SC	II	S
Savanna Hawk	<i>Buteogallus meridionalis</i>	3,9	SC	II	S
Harris's Hawk	<i>Parabuteo unicinctus</i>	6,7	SC	III	S
Roadside Hawk	<i>Buteo magnirostris</i>	1–9,12,13,15–18	C	II,III	R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Broad-winged Hawk	<i>Buteo platypterus</i>	1	X	II	S,H
Red-backed Hawk	<i>Buteo polyosoma</i>	1–3,5,8,9	U	II,III,V	R,P,S,H
Short-tailed Hawk	<i>Buteo brachyurus</i>	13	X	II	S
Zone-tailed Hawk	<i>Buteo albonotatus</i>	8	X	II	S
Black Hawk-Eagle	<i>Spizaetus tyrannus</i>	3–9,17,18	FC	II,V	R,P,S,H
Ornate Hawk-Eagle	<i>Spizaetus ornatus</i>	3	X	II	S
Falconidae					
Laughing Falcon	<i>Herpetotheres cachinnans</i>	1,3–10,13,18	FC	II,III	R,S,H
Barred Forest-Falcon	<i>Micrastur ruficollis</i>	1–5,7–9,18	FC	I,II	R,S,H
Crested Caracara	<i>Caracara cheriway</i>	1–4,11,13,17,18	U	III	S
Red-throated Caracara	<i>Ibycter americanus</i>	3,5–9,11,13,15–18	FC	I,II	R,S,H
Yellow-headed Caracara	<i>Milvago chimachima</i>	1,2,8,9,13	U	II,III	S,H
American Kestrel	<i>Falco sparverius</i>	1,3–11,13,14,16–18	C	III	R,S,H
Bat Falcon	<i>Falco rufigularis</i>	18	SC	II	S,H
Aplomado Falcon	<i>Falco femoralis</i>	2	X	III	S
Rallidae					
Russet-crowned Crake	<i>Anurolimnas viridis</i>	1,4,5,7,8,13,14	FC	III	R,S,H
Charadriidae					
Southern Lapwing	<i>Vanellus chilensis</i>	1,3,5,8,9,13,18	U	III	S,H
Columbidae					
Ruddy Ground-Dove	<i>Columbina talpacoti</i>	8,9,18	U	III	S
Scaled Pigeon	<i>Patagioenas speciosa</i>	6,8,16,18	U	II	R,S,H
Band-tailed Pigeon	<i>Patagioenas fasciata</i>	1–9,11,13,16,18	C	I,II,III	R,S,H
Pale-vented Pigeon	<i>Patagioenas cayennensis</i>	4,7–9,18	U	II,III	S,H
Ruddy Pigeon	<i>Patagioenas subvinacea</i>	1–10,13,17,18	C	I,II	R,S,H
Ground-Dove	<i>Geotrygon m. montana</i>	2,6,7,10	SC	I	M,R,H
Eared Dove	<i>Zenaidura auriculata</i>	2,8,13	SC	III	S,H
White-tipped Dove	<i>Leptotila verreauxi</i>	8,9	SC	II	S
Psittacidae					
Blue-fronted Parrotlet	<i>Touit dilectissimus</i>	1–9,11–14,16–18	C	I,II	R,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Saffron-headed Parrot (NT)	<i>Pyrilia pyrilia</i>	1–3,5–8,13,17,18	FC	I,II	M,R,P,S,H
Blue-headed Parrot	<i>Pionus menstruus</i>	7,14	SC	I,II	S,H
Bronze-winged Parrot	<i>Pionus c. chalcopterus</i>	4,5,7,8,11,13,14,17,18	FC	I,II	R,P,S,H
Mealy Parrot	<i>Amazona farinosa</i>	17,18	U	I,II	R,P,S,H
Cuculidae					
Squirrel Cuckoo	<i>Piaya cayana</i>	1–9,15,16,18	FC	I,II,III	R,P,S,H
Smooth-billed Ani	<i>Crotophaga ani</i>	1–11,13,14,16–18	C	II,III	R,S,H
Striped Cuckoo	<i>Tapera naevia</i>	1,2,4,5,7–9,12,13,16,17	FC	II,III	R,H
Pavonine Cuckoo	<i>Dromococcyx pavoninus</i>	3,8,9,15,18	U	II,III	R,H
Strigidae					
Tropical Screech-Owl	<i>Megascops choliba</i>	1–5,7–10,13,18	FC	II,III	R,S,H
Cinnamon Screech-Owl	<i>Megascops petersoni</i>	8,9	U	I,II	M,R,P,H
Mottled Owl	<i>Ciccaba virgata</i>	1–4,7–9,13	FC	I,II	R,S,H
Nyctibiidae					
Common Potoo	<i>Nyctibius griseus</i>	3,4,7–9,13	FC	II	R,P,S,H
Caprimulgidae					
Common Nighthawk ⁶	<i>Chordeles minor</i>	7	SC	II	S
Common Pauraque	<i>Nyctidromus albicollis</i>	2–4,6–10,13,14,16	C	III	S,H
Band-winged Nightjar	<i>Caprimulgus longirostris ruficervix</i>	1–3,5–9,16,18	C	II,III	M,R,P,S,H
Apodidae					
Chestnut-collared Swift	<i>Streptoprocne rutila</i>	1–10,13,16–18	C	V	R,P,S,H
White-collard Swift	<i>Streptoprocne zonaris</i>	1–4,7–10,13,15–18	FC	V	P,S,H
Gray-rumped Swift	<i>Chaetura cinereiventris</i>	3,7,8	FC	V	R,S,H
Short-tailed Swift	<i>Chaetura brachyura</i>	13	X	V	S
Lesser Swallow-tailed Swift	<i>Panyptila cayennensis</i>	7	SC	V	S
Trochilidae					
White-necked Jacobin	<i>Florisuga mellivora</i>	3,13	SC	II,III	S
White-tipped Sicklebill	<i>Eutoxeres aquila munda</i>	1–5,7–10,18	C	I,II	M,P,S
Rufous-breasted Hermit	<i>Glaucis hirsutus affinis</i>	13	SC	II	M,P
Band-tailed Barbetthroat	<i>Threnetes ruckeri darienensis</i>	10	SC	II	M

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Stripe-throated Hermit	<i>Phaethornis s. strigularis</i>	4,7,8	U	I,II	M,P,S
Green Hermit	<i>Phaethornis guy emiliae</i>	1–10,13–16,18	C	I,II	M,R,P,S,H
Tawny-bellied Hermit	<i>Phaethornis s. gyrmathopborus</i>	8	SC	II	M,P,S
Green-fronted Lancebill	<i>Doryfera l. ludoviciae</i>	1,3–5,7–10,13,16,18	C	I,II	M,P,S
Wedge-billed Hummingbird	<i>Schistes geoffroyi albogularis</i>	2,3	U	I,II	M,P,S
Brown Violetear	<i>Colibri delphinae</i>	3	X	I	M,P
Green Violetear	<i>Colibri thalassinus</i>	1	X	II	S,H
Sparkling Violetear	<i>Colibri c. coruscans</i>	1	U	II,III	M,P,S,H
Tooth-billed Hummingbird	<i>Androdon aequatorialis</i>	7,18	U	I	M,P,S,H
Purple-crowned Fairy	<i>Heliothrix barroti</i>	7	X	I	S
Black-throated Mango	<i>Anthracothorax nigricollis</i>	1,4,7	SC	III	S
Greenish Puffleg	<i>Haplophaedia a. aureliae</i>	1–3,7–10,12,15,16,18	C	I,II	M,R,P,S,H
Bronzy Inca	<i>Coeligena coeligena ferruginea</i>	1–3,8,9,15,16	FC	I,II	M,R,P,S,H
Booted Racket-tail	<i>Oreotrochilus underwoodii</i>	1–3,5–9,15,16,18	C	I,II	M,R,P,S,H
Green-crowned Brilliant	<i>Heliodoxa j. jacula</i>	1–10,13,16,18	C	I,II	M,R,P,S,H
Purple-throated Woodstar	<i>Calliphlox mitchelli</i>	5,9	SC	I,II	M,P,S
Western Emerald	<i>Chlorostilbon melanorhynchus pumilus</i>	1–5,7,9–11,13–15,17	C	I,II,III	M,P,S
Violet-headed Hummingbird	<i>Klais guimeti</i>	7,18	SC	I	M,P
Green-crowned Woodnymph	<i>Thalurania f. fannyi</i>	3–10,12,13,16,18	C	I,II,III	M,P,S
Rufous-tailed Hummingbird	<i>Amazilia t. tzacatl</i>	4,6–8,11,13	U	III	R,S,H
Andean Emerald	<i>Amazilia f. franciae</i>	1–9,11,13–18	C	I,II	M,R,P,S,H
Steely-vented Hummingbird	<i>Amazilia s. saucerottei</i>	1–5,7,8,13	FC	II,III	M,P,S
Trogonidae					
Golden-headed Quetzal	<i>Pharomachrus auriceps</i>	1–4,7–10,13,15–18	FC	I,II	M,R,P,S,H
Collared Trogon	<i>Trogon collaris subtropica</i>	1–10,13–18	FC	I,II	M,R,P,S,H
Blue-tailed Trogon	<i>Trogon comptus</i>	18	X	I	R,S,H
Alcedinidae					
Green Kingfisher	<i>Chloroceryle a. americana</i>	9,13,17,18	SC	II,IV	M,P,S
Momotidae					
Blue-crowned Motmot	<i>Momotus momota aequatorialis</i>	1–4,7–9	U	I,II,III	P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Bucconidae					
Moustached Puffbird	<i>Malacoptila m. mystacalis</i>	1–10,13,16–18	FC	I,II	M,R,P,S,H
Lanceolated Monklet	<i>Micromonacha lanceolata</i>	7,8	SC	I,II	S,H
Capitonidae					
White-mantled Barbet (EN)	<i>Capito b. hypoleucus</i>	2,7,8,10	U	I,II	M,R,P,S,H
Red-headed Barbet	<i>Eubucco b. bourcieri</i>	1–10,16,18	FC	I,II	M,R,P,S,H
Ramphastidae					
Black-mandibled Toucan	<i>Ramphastos ambiguus swainsonii</i>	7,9,17,18	SC	I,II,III	R,H
Channel-billed Toucan	<i>Ramphastos vitellinus citrolaemus</i>	3	X	II	S
Crimson-rumped Toucanet	<i>Aulacorhynchus h. haematopygus</i>	1–5,7–10	FC	I,II	M,R,P,S,H
Picidae					
Olivaceous Piculet	<i>Picumnus olivaceus</i>	1–4,7,9,13,15	U	I,II,III	M,P,S
Acorn Woodpecker	<i>Melanerpes formicivorus flavigula</i>	1–3,5,7–9,11,13,14,16,18	U	II,III	M,R,P,S,H
Red-crowned Woodpecker	<i>Melanerpes rubricapillus</i>	1,8–10	SC	III	S,H
Smoky-brown Woodpecker	<i>Picoides f. fumigatus</i>	1–10,13,16,18	FC	I,II	M,R,P,S,H
Golden-olive Woodpecker	<i>Colaptes rubiginosus gularis</i>	1–5–8,10,18	FC	I,II,III	R,P,S,H
Spot-breasted Woodpecker	<i>Colaptes punctigula</i>	1,4–6,8	U	III	S,H
Lineated Woodpecker	<i>Dryocopus lineatus</i>	3–5,7,8,11,13,14,18	FC	II,III	P,S,H
Crimson-crested Woodpecker	<i>Campetherus melanoleucus</i>	1–5,7–9,13	FC	I,II,III	R,P,S,H
Furnariidae					
Tawny-throated Leafcutter	<i>Sclerurus mexicanus andinus</i>	7	SC	I	M,P,H
Azara's Spinetail	<i>Synallaxis azarae media</i>	1–9,11,12,14,16–18	C	I,II,III	M,R,P,S,H
Pale-breasted Spinetail	<i>Synallaxis albescens</i>	1–9,11,13,14	FC	II,III	R,P,S,H
Slaty Spinetail	<i>Synallaxis brachyura</i>	7	SC	II	S
Red-faced Spinetail	<i>Cranioleuca erythrops griseigularis</i>	1–3,5,7–9,18	FC	I,II	M,R,P,S,H
Spotted Barbtail	<i>Premnoplex b. brunneiceps</i>	2,3,5,7–9,18	FC	I	M,R,P,S,H
Treerunner	<i>Margarornis</i> sp.	7	X	II	S
Montane Foliage-gleaner	<i>Anabacerthia striaticollis</i>	2,3	SC	I	P,S
Lineated Foliage-gleaner	<i>Syndactyla subalaris striolata</i>	1–5,7–9,16,18	FC	I,II	M,R,P,S,H
Striped Woodhaunter	<i>Hyloctistes subulatus cordobae</i>	5–7,9,17,18	FC	I	M,R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Slaty-winged Foliage-gleaner	<i>Philydor fuscipenne erythronotum</i>	7,18	U	I	M,R,P,S,H
Buff-fronted Foliage-gleaner	<i>Philydor rufum riveti</i>	1–9,12,13,15–18	C	I,II	M,R,P,S,H
Streak-capped Treecrusher	<i>Thripadectes virgaticeps magdalena</i>	1–3,5,7–9,15	U	I	M,P,S,H
Ruddy Foliage-gleaner	<i>Automolus rubiginosus saturatus</i>	5,7,18	SC	I	M,R,P,S,H
Plain Xenops	<i>Xenops minutus littoralis</i>	4,5,7–9,18	U	I,II	M,R,P,S,H
Streaked Xenops	<i>Xenops rutilans heterurus</i>	1–5,7,8,13,15,18	FC	I,II	M,R,P,S,H
Plain-brown Woodcreeper	<i>Dendrocincla fuliginosa</i>	7,9,18	SC	I	P,S,H
Olivaceous Woodcreeper	<i>Sittasomus griseicapillus levis</i>	2–5,7–10,15,17,18	FC	I,II	M,R,P,S,H
Wedge-billed Woodcreeper	<i>Glyphorynchus spirurus integratus</i>	7	SC	I	M,P,S
Strong-billed Woodcreeper	<i>Xiphocolaptes promeropirhynchus</i>	3,5,7,9	U	I	S,H
Black-banded Woodcreeper	<i>Dendrocolaptes picumnus</i>	9	SC	I	R,H
Spotted Woodcreeper	<i>Xiphorhynchus erythropygius insolitus</i>	4–7,10,13,18	FC	I	M,R,P,S,H
Olive-backed Woodcreeper	<i>Xiphorhynchus t. triangularis</i>	1–3,8,9,16	U	I	M,P,S,H
Montane Woodcreeper	<i>Lepidocolaptes lacrymiger</i>	2,3,5,9	U	I,II	P,S,H
Brown-billed Scythebill	<i>Campylorhamphus p. pusillus</i>	2,3,5,7–9,16,18	FC	I	M,R,P,S,H
Thamnophilidae					
Bar-crested Antshrike	<i>Thamnophilus multistriatus</i>	1,3–5,7–9,12	FC	II,III	R,S,H
Uniform Antshrike	<i>Thamnophilus unicolor grandior</i>	1–4,6–9,15–18	FC	I,II	M,R,P,S,H
Plain Antvireo	<i>Dysithamnus mentalis semicinereus</i>	2–5,7–10,18	FC	I,II	M,R,P,S,H
Checker-throated Antwren	<i>Epinecrophylla fulviventris salmoni</i>	5	X	I	M,P
Slaty Antwren	<i>Myrmotherula s. schisticolor</i>	2–10,18	FC	I	M,R,P,S,H
Long-tailed Antbird	<i>Drymophila c. caudata</i>	1,2	U	I,II	M,R,P,S,H
Rufous-rumped Antwren	<i>Terenura c. callinota</i>	2,3,7–9,18	U	I,II	M,R,S,H
Parker's Antbird	<i>Cercomacra parkeri</i>	1–10,13,15–18	C	I,II,III	M,R,P,S,H
Immaculate Antbird	<i>Myrmeciza i. immaculata</i>	7,15,18	SC	I	R,S,H
Formicariidae					
Rufous-breasted Anthrush	<i>Formicarius rufippectus carrikeri</i>	3,9,10	SC	I	M,R,S,H
Grallariidae					
White-bellied Antpitta	<i>Grallaria b. hypoleuca</i>	2,3,8,9,16	U	I	R,S,H
Ochre-breasted Antpitta	<i>Grallaricula flavirostris</i>	2,5,8,9,16	U	I	M,R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Conopophagidae					
Chestnut-crowned Gnat-eater	<i>Conopophaga c. castaneiceps</i>	4–7,9,10,13,18	U	I	M,R,P,S,H
Rhinocryptidae					
White-crowned Tapaculo	<i>Scytalopus atratus confusus</i>	3–5,7–9,13,18	FC	II	M,R,P,S,H
Stiles's Tapaculo (NT)	<i>Scytalopus stilesi</i>	1–10,13,15–18	C	I	M,R,P,S,H
Tyrannidae					
Sooty-headed Tyrannulet	<i>Phyllomyias griseiceps</i>	7,8,17	U	II	R,S,H
Ashy-headed Tyrannulet	<i>Phyllomyias cinereiceps</i>	2	SC	I,II	M,P
Plumbeous-crowned Tyrannulet	<i>Phyllomyias plumbeiceps</i>	2,3,5,7	U	I	M,R,P,S,H
Gray Elaenia	<i>Myiopagis caniceps</i>	3,7,9	U	I	M,R,P,S,H
Yellow-bellied Elaenia	<i>Elaenia flavogaster</i>	1,2,4,5,7–9,13,15	FC	II,III	R,P,S,H
Lesser Elaenia	<i>Elaenia chiriquensis albivertex</i>	13	SC	II	M,P
Mountain Elaenia	<i>Elaenia frantzii</i>	1,2,5,7	U	II	R,P,S,H
Torrent Tyrannulet	<i>Serpophaga cinerea</i>	5–7,9	FC	IV	S
Golden-faced Tyrannulet	<i>Zimmerius c. chrysops</i>	1–9,11–18	C	I,II,III	M,R,P,S,H
Marble-faced Bristle-Tyrant	<i>Phylloscartes o. opthalmicus</i>	1–3,5,7–9,13,17,18	FC	I,II	M,R,P,S,H
Rufous-browed Tyrannulet	<i>Phylloscartes superciliosus</i>	1–5,7–9,13–15,17,18	FC	I,II	M,R,P,S,H
Streak-necked Flycatcher	<i>Mionectes striaticollis columbianus</i>	1–3,8,10,15,16	C	I,II	M,R,P,S,H
Olive-striped Flycatcher	<i>Mionectes olivaceus hederaceus</i>	1,4,6–9,13,15,16,18	C	I,II	M,P,S
Ochre-bellied Flycatcher	<i>Mionectes oleagineus</i>	4,7,11	SC	I,II	P,S,H
Slaty-capped Flycatcher	<i>Leptopogon superciliaris poliocephalus</i>	1,3–5,7,8,13,15,18	FC	I,II,III	M,R,P,S,H
Ornate Flycatcher	<i>Myiobius ornatus stellatus</i>	2–9,12,13,16–18	FC	I,II	M,R,P,S,H
Scale-crested Pygmy-Tyrant	<i>Lophotriccus pileatus hesperius</i>	1–11,13–18	C	I,II	M,R,P,S,H
Pearly-vented Tody-Tyrant	<i>Hemitriccus margaritaceiventer</i>	3,10	SC	II	R,S,H
Common Tody-Flycatcher	<i>Todirostrum cinereum</i>	4,5,7–9,11,13,14,17	SC	III	S,H
Fulvous-breasted Flatbill	<i>Rhynchocyclus fulvipectus</i>	2,5,7,9	SC	I	M,P,S
White-throated Spadebill	<i>Platyrinchus mystaceus neglectus</i>	2–9,18	FC	I	M,R,P,S,H
Orange-crested Flycatcher	<i>Myioborus phoenicomitra latae</i>	7	SC	I	M,P,S
Tawny-breasted Flycatcher	<i>Myiobius v. villosus</i>	5,7,18	SC	I	M,P,S
Cinnamon Flycatcher	<i>Pyrrhomyias cinnamomeus pyrrhopterus</i>	1–3,8,9,17	U	I,II	M,R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Acadian Flycatcher ⁶	<i>Empidonax virescens</i>	4,7,8,13,18	FC	I,II,III	M,P,S
Willow Flycatcher ⁶	<i>Empidonax traillii</i>	13	SC	I,II,III	M,P,S
Alder Flycatcher ⁶	<i>Empidonax alnorum</i>	2–5,7	SC	I,II,III	M,P,S
Olive-sided Flycatcher ⁶ (NT)	<i>Contopus cooperi</i>	3,6–9	SC	I,II,III	R,S,H
Smoke-colored Pewee	<i>Contopus fumigatus ardosiacus</i>	2,3,7,9,16	U	I,II	M,R,P,S,H
Eastern Wood-Pewee ⁶	<i>Contopus virens</i>	3–8,13,17,18	FC	II,III	R,P,S,H
Western Wood-Pewee ⁶	<i>Contopus sordidulus</i>	7	SC	II,III	R,H
Black Phoebe	<i>Sayornis nigricans</i>	1,4,6–9,13,15,17,18	C	3,4	S,H
Vermillion Flycatcher	<i>Pyrocephalus rubinus</i>	4	SC	III	S
Rufous-tailed Tyrant	<i>Knipolegus poecilurus</i>	3	SC	II	R,P,S,H
Cattle Tyrant	<i>Machaetornis rixosa</i>	7–9,11	FC	III	S,H
Piratic Flycatcher	<i>Legatus leucophaius</i>	5,7,8	SC	II	H
Rusty-margined Flycatcher	<i>Myiozetetes cayanensis</i>	3–10,13,14	C	III	R,P,S,H
Great Kiskadee	<i>Pitangus sulphuratus</i>	1,4,8,9	U	III	R,S,H
Lemon-browed Flycatcher	<i>Conopias cinchoneti</i>	18	X	II	S
Golden-crowned Flycatcher	<i>Myiodynastes chrysocephalus minor</i>	1–9,13,18	FC	I,II	R,P,S,H
Streaked Flycatcher	<i>Myiodynastes maculatus</i>	5,7–9	U	II,III	R,S,H
Fork-tailed Flycatcher	<i>Tyrannus savanna</i>	1–9,11,13	FC	III	S,H
Tropical Kingbird	<i>Tyrannus melancholicus</i>	1–16,18	C	II,III	R,P,S,H
Rufous Mourner	<i>Rhytipterna holerythra</i>	7	SC	I	S
Dusky-capped Flycatcher	<i>Myiarchus t. tuberculifer</i>	1–5,7–9,13,18	FC	I,II	M,R,P,S,H
Pale-edged Flycatcher	<i>Myiarchus cephalotes canace</i>	1–3,5–9,13,16–18	FC	I,II	M,R,P,S,H
Great Crested Flycatcher ⁵	<i>Myiarchus crinitus</i>	13	X	II	M,P,S
Bright-rumped Attila	<i>Attila spadiceus sclateri</i>	2–5,7,8	FC	I,II	M,R,P,S,H
Oxyruncidae					
Sharpbill	<i>Oxyruncus cristatus</i>	18	X	I	M,P
Cotingidae					
Golden-breasted Fruit-eater	<i>Pipreola a. aureopectus</i>	2,3,8,9	U	I,II	M,R,P,S,H
Chestnut-capped Piha (EN)	<i>Lipaugus weberi</i>	2–9,11–18	FC	I,II	M,R,P,S,H
Purple-throated Fruitcrow	<i>Querula purpurata</i>	2,3,5,7,18	U	I,II	M,R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Red-ruffed Fruitcrow	<i>Pyroderus scutatus occidentalis</i>	2,3,9	U	I,II	R,P,S,H
Pipridae					
Golden-winged Manakin	<i>Masius chrysopterus bellus</i>	1–3,5,7–9,11,16	FC	I,II	M,R,P,S,H
Striped Manakin	<i>Machaeropterus regulus antioquiae</i>	4,7,9,10,18	U	I,II	M,R,P,S,H
Blue-crowned Manakin	<i>Lepidothrix coronata</i>	18	X	II	S
White-bearded Manakin	<i>Manacus manacus abditivus</i>	2,9	SC	II	M,P,H
Green Manakin	<i>Xenopipo holochlora litiae</i>	18	X	I	M,P
Yellow-headed Manakin (NT)	<i>Xenopipo flavicapilla</i>	2	U	I	M,P
White-crowned Manakin	<i>Pipra pipra bolivari</i>	4–7,9,10,13,18	C	I,II	M,R,P,S,H
Golden-headed Manakin	<i>Pipra erythrocephala</i>	5,18	X	I	P
Tityridae					
Masked Tityra	<i>Tityra semifasciata columbianus</i>	7,9,10	SC	II	P,S,H
Thrush-like Schiffornis	<i>Schiffornis turdina panamensis</i>	18	X	I	M,P,H
Barred Becard	<i>Pachyramphus v. versicolor</i>	2,3,9,10,16–18	U	I,II	R,P,S,H
Cinereous Becard	<i>Pachyramphus r. rufus</i>	7,13	U	II	M,P,S
Cinnamon Becard	<i>Pachyramphus cinnamomeus</i>	3,4,6,7,10,18	FC	II,III	R,S,H
White-winged Becard	<i>Pachyramphus polychopterus cinereiventris</i>	1–8,16,18	FC	I,II	M,R,P,S,H
Incuba Sedis 1					
Wing-barred Piprites	<i>Piprites chloris antioquiae</i>	2,3,5–8,16–18	FC	I,II	M,R,P,S,H
Vireonidae					
Black-billed Peppershrike	<i>Clytarpus nigrirostris</i>	1–5,7–9	FC	I,II	R,P,S,H
Yellow-browed Shrike-Vireo	<i>Vireolanius eximius</i>	7,10	U	I	R,S,H
Yellow-throated Vireo ⁶	<i>Vireo flavifrons</i>	5	X	II	S
Brown-capped Vireo	<i>Vireo l. leucophrys</i>	1–3,8,18	FC	I,II	M,R,P,S,H
Red-eyed Vireo ⁶	<i>Vireo olivaceus</i>	2,8,13,18	SC	II	R,P,S,H
Rufous-naped Greenlet	<i>Hylophilus semibrunneus</i>	1–9,13,18	C	I,II	M,R,P,S,H
Corvidae					
Black-chested Jay	<i>Cyanocorax affinis</i>	1,2,4,5,8–10,18	U	II,III	R,S,H
Green Jay	<i>Cyanocorax yncas galeatus</i>	1–10,13,16–18	C	I,II	M,R,P,S,H
Hirundinidae					

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Blue-and-white Swallow	<i>Pygochelidon cyanoleuca</i>	1–9,11–13,16–18	FC	III,V	S,H
White-thighed Swallow	<i>Atticora tibialis minima</i>	6–8	U	II,V	M,R,P,S,H
Southern Rough-winged Swallow	<i>Stelgidopterix ruficollis</i>	1–18	C	III,V	R,P,S,H
Gray-breasted Martin	<i>Progne chalybea</i>	7	U	IS,V	S
Troglodytidae					
Scaly-breasted Wren	<i>Microcerculus marginatus</i>	7,9,10	U	I	R,S,H
House Wren	<i>Troglodytes aedon striatulus</i>	1–9,11–14,17,18	FC	II,III	M,R,P,S,H
Sooty-headed Wren	<i>Thryothorus spadix</i>	1–10,13,15–18	C	I,II	M,R,P,S,H
Whiskered Wren	<i>Thryothorus mystacalis</i>	1	SC	II,III	H
Bay Wren	<i>Thryothorus nigricapillus</i>	5–7,9,15,18	U	II,IV	R,S,H
Gray-breasted Wood-Wren	<i>Henicorhina l. leucophrys</i>	2–18	FC	I,II	M,R,P,S,H
Song Wren	<i>Cyphorhinus phaeocephalus</i>	7	X	I	H
Polioptilidae					
Long-billed Gnatwren	<i>Ramphocaenus melanurus griseodorsalis</i>	1,7,9	U	I,II	M,R,P,S,H
Turdidae					
Andean Solitaire	<i>Myadestes ralloides plumbeiceps</i>	1–3,6–9,16,18	FC	I,II	M,R,P,S,H
Gray-cheeked Thrush ⁶	<i>Catharus minimus</i>	5,7–9,13,16,18	U	I,II	P,S,H
Swainson's Thrush ⁶	<i>Catharus ustulatus</i>	1–5,7–10,13,15,18	C	I,II,III	R,P,S,H
Pale-eyed Thrush	<i>Turdus leucomelas</i>	3,5,7–9,13,15,18	FC	I,II	M,R,P,S,H
Great Thrush	<i>Turdus fuscater quindio</i>	1–3,6,8,9,11,12,14–17	FC	II,III	M,R,P,S,H
Glossy-black Thrush	<i>Turdus serranus</i>	2,3,9,16	SC	I	S
Black-billed Thrush	<i>Turdus i. ignobilis</i>	1–9,13	FC	II,III	R,P,S,H
Pale-vented Thrush	<i>Turdus o. obsoletus</i>	7	SC	I,II	M,P,S
Mimidae					
Tropical Mockingbird	<i>Mimus gilvus tolimensis</i>	1–9,13	FC	III	M,R,P,S,H
Thraupidae					
Black-faced Tanager	<i>Schistochlamys melanopis</i>	1,3,6–9,12–14,17,18	U	II,III	S
Oleaginous Hemispingus	<i>Hemispingus f. frontalis</i>	2,3,8,9	FC	I,II	M,R,P,S,H
White-lined Tanager	<i>Tachyphonus rufus</i>	1,2,5	U	III	S,H
Crimson-backed Tanager	<i>Ramphocelus dimidiatus</i>	1–10,12,13,15,17	FC	III	S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Flame-rumped Tanager	<i>Ramphocelus flammigerus icteronotus</i>	4,6–14,17,18	FC	III	S,H
Blue-gray Tanager	<i>Thraupis episcopus canus</i>	1–14,17,18	FC	II,III	R,P,S,H
Palm Tanager	<i>Thraupis palmarum atripennis</i>	1,3–5,7–11,13,17,18	FC	II,III	R,P,S,H
Black-and-gold Tanager (VU)	<i>Bangsia melanochlamys</i>	8,9,16–18	U	I,II	M,R,P,S,H
Blue-winged Mountain-Tanager	<i>Anisognathus somptuosus antioquiae</i>	1–4,6,8,9,18	FC	I,II,III	M,R,P,S,H
Purplish-mantled Tanager (NT)	<i>Iridosornis porphyrocephalus</i>	1–3,6–9,15–18	FC	I,II,III	M,R,P,S,H
Multicolored Tanager (VU)	<i>Chlorocrysa nitidissima</i>	2,3,5,7–9	U	I,II	M,R,P,S,H
Golden Tanager	<i>Tangara arthus occidentalis</i>	1–5,7–9,11–18	C	I,II	M,R,P,S,H
Silver-throated Tanager	<i>Tangara i. icterocephala</i>	2,3,7,8	FC	I,II	M,R,P,S,H
Saffron-crowned Tanager	<i>Tangara xanthocephala</i>	3	X	II	S
Speckled Tanager	<i>Tangara guttata tolimae</i>	3–10,13,14,18	U	I,II	M,R,P,S,H
Bay-headed Tanager	<i>Tangara gyrola deleticia</i>	1,3–5,7–9,13,14,17,18	FC	I,II,III	M,R,P,S,H
Scrub Tanager	<i>Tangara vitriolina</i>	1–9,11–14,16,17	FC	II,III	R,P,S,H
Blue-necked Tanager	<i>Tangara cyanicollis granadensis</i>	1–5,7–9,11–13,14,17	U	II,III	M,R,P,S,H
Blue-necked Tanager	<i>Tangara nigroviridis consobrina</i>	1–9,11,13–18	FC	I,II	M,R,P,S,H
Black-capped Tanager	<i>Tangara heinei</i>	1–5,7,8,11,13,16–18	FC	II,III	M,R,P,S,H
Purple Honeycreeper	<i>Cyanerpes caeruleus microrhynchus</i>	7,18	SC	I	M,P,S
Green Honeycreeper	<i>Chlorophanes spiza</i>	9	SC	I,II	S
Guira Tanager	<i>Hemithraupis guira guirina</i>	2–5,7–9,13,18	U	I,II	M,P,S
Scarlet-and-white Tanager	<i>Chrysotthlypis salmoni</i>	7,15,18	FC	I,II	M,R,P,S,H
Rusty Flowerpiercer	<i>Diglossa sittoides dorbignyi</i>	1–3	U	II,III	M,R,P,S,H
Indigo Flowerpiercer	<i>Diglossa indigotica</i>	8,9,16–18	SC	I,II	M,R,P,S,H
Incerta Sedis 2					
Common Bush-Tanager	<i>Chlorospingus opthalmicus nigriceps</i>	1,2	U	II	P,S
Yellow-throated Bush-Tanager	<i>Chlorospingus f. flavigularis</i>	2–4,7–10,13,15–18	C	I,II	M,R,P,S,H
Bananaquit	<i>Coereba flaveola columbiana</i>	1–9,11,13,15–18	C	I,II,III	R,P,S,H
Yellow-faced Grassquit	<i>Tiaris olivaceus</i>	1–13,17	C	II,III	R,P,S,H
Dull-colored Grassquit	<i>Tiaris o. obscurus</i>	5	SC	II	M,P,S
Buff-throated Saltator	<i>Saltator maximus</i>	4,5,7–10,13	FC	II,III	R,P,S,H
Black-winged Saltator	<i>Saltator a. atripennis</i>	1–9,12,13,17,18	FC	I,II	M,R,P,S,H

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Grayish Saltator	<i>Saltator coerulescens</i>	7–9	U	II,III	S
Streaked Saltator	<i>Saltator striaticeps</i>	1,4,5,9	FC	II,III	R,S,H
Emberizidae					
Rufous-collared Sparrow	<i>Zonotrichia capensis costaricensis</i>	1–9,11–14,16–18	C	II,III	M,R,P,S,H
Slaty Finch	<i>Haplospiza r. rustica</i>	15	SC	I	M,R,P,S,H
Saffron Finch	<i>Sicalis flaveola</i>	6–8	SC	III	S,H
Blue-black Grassquit	<i>Volatinia jacarina</i>	3–5,7–9,13,14	U	III	S,H
Slate-colored Seedeater	<i>Sporophila schistacea</i>	9,16	SC	II	S
Variable Seedeater	<i>Sporophila corvina</i>	10	X	III	S
Yellow-bellied Seedeater	<i>Sporophila nigricollis</i>	1,3–9,13	C	II,III	R,P,S,H
Ruddy-breasted Seedeater	<i>Sporophila minuta</i>	7–9,11,13,17,18	U	III	S
Thick-billed Seed-Finch	<i>Oryzoborus funereus ochrogynne</i>	5,7,10,13,15	U	I	M,P,S
Black-striped Sparrow	<i>Arremonops conirostris</i>	4–9,13,17,18	FC	III	R,S,H
Chestnut-capped Brush-Finch	<i>Arremon brunneinucha frontalis</i>	1–9,11,13,17,18	FC	I,II	M,R,P,S,H
Stripe-headed Brush-Finch	<i>Arremon torquatus atricapillus</i>	5,7,9	U	II	M,R,P,S,H
White-naped Brush-Finch	<i>Atlapetes albiniucha gutturalis</i>	1–9,13,15,16,18	FC	II,III	M,R,P,S,H
Cardinalidae					
Summer Tanager ⁶	<i>Piranga rubra</i>	1–5,6,8,9,12,13,17,18	FC	I,II,III	P,S,H
Scarlet Tanager ⁶	<i>Piranga olivacea</i>	8,13,18	FC	II	P
White-winged Tanager	<i>Piranga leucoptera venezuelae</i>	2,3,5,7–9,15,18	U	I,II	M,R,P,S,H
Sooty Ant-Tanager (NT)	<i>Habia gutturalis</i>	7	X	I	M,R,S,H
Crested Ant-Tanager	<i>Habia cristata</i>	18	SC	I,IV	R,S,H
Lemon-spectacled Tanager	<i>Chlorothraupis olivacea</i>	7	X	I	S
Rose-breasted Grosbeak ⁶	<i>Pheucticus ludovicianus</i>	2–5,13,18	U	II,III	P,S
Blue-black Grosbeak	<i>Cyanocompsa c. cyanoides</i>	5,7,10	U	I,II	M,R,P,S,H
Parulidae					
Golden-winged Warbler ⁶ (NT)	<i>Vermivora chrysoptera</i>	4,5,8,9,16	U	I	S
Tennessee Warbler ⁶	<i>Vermivora peregina</i>	3,5,8	U	I,II	P,S
Tropical Parula	<i>Parula pityayumi</i>	7,9	U	II,III	S
Bay-breasted Warbler ⁶	<i>Dendroica castanea</i>	2,4,5,7,13,17	FC	II	P,S

APPENDIX 1. Continued.

English names ¹	Scientific names ²	Localities	Abundance ³	Habitats ⁴	Evidence ⁵
Blackburnian Warbler ⁶	<i>Dendroica fusca</i>	1–9,11–14,16–18	C	I,II,III	P,S
Yellow Warbler ⁶	<i>Dendroica petechia</i>	6	U	II	S
American Redstart ⁶	<i>Setophaga ruticilla</i>	4	U	II	P,S
Black-and-white Warbler ⁶	<i>Minotilla varia</i>	1–5,7–9,11,16,18	FC	I,II	S
Northern Waterthrush ⁶	<i>Seiurus noveboracensis</i>	4,13	U	II,IV	P,S
Mourning Warbler ⁶	<i>Oporornis philadelphica</i>	4,13,18	U	II	P,S
Canada Warbler ⁶	<i>Wilsonia canadensis</i>	1–5,7,8,10,13,18	FC	I,II	P,S
Slate-throated Redstart	<i>Myiothorus miniatus ballux</i>	1–9,13,15–18	C	I,II,III	M,R,P,S,H
Golden-crowned Warbler	<i>Basileuterus culicivorus</i>	7	SC	I,II	P,S
Rufous-capped Warbler	<i>Basileuterus rufifrons mesochrysus</i>	3,4	U	II,III	M,P,S
Three-striped Warbler	<i>Basileuterus tristriatus</i>	1–5,7–12,15–18	C	I,II	M,R,P,S,H
Buff-rumped Warbler	<i>Phaeothlypis fulvicauda</i>	6,7,18	U	II,IV	R,S,H
Icteridae					
Russet-backed Oropendola	<i>Psarocolius angustifrons salmoni</i>	1–10,13,15–18	C	I,II,III	R,P,S,H
Scarlet-rumped Cacique	<i>Cacicus uropygialis</i>	7,18	SC	I,II	R,S,H
Red-bellied Grackle (EN)	<i>Hypopyrrhus pyrohypogaster</i>	1–10,12,13,15–18	C	I,II,III	M,R,P,S,H
Giant Cowbird	<i>Molothrus oryzivorus</i>	3,4,7,9	SC	III	S
Shiny Cowbird	<i>Molothrus bonariensis</i>	3,4,7,8,12,13	U	III	S,H
Fringillidae					
Andean Siskin	<i>Carduelis spinescens</i>	8	X	II	S
Yellow-bellied Siskin	<i>Carduelis x. xanthogastra</i>	2,3,7–9	U	II	M,R,P,S,H
Lesser Goldfinch	<i>Carduelis psaltria</i>	1,4–9,13,16	FC	III	S,H
Thick-billed Euphonia	<i>Euphonia laniirostris</i>	4,8,9	U	III	S,H
Golden-rumped Euphonia	<i>Euphonia cyanocephala pelzelni</i>	5,7,13	SC	II	M,P,S
Orange-bellied Euphonia	<i>Euphonia xanthogaster oreassinoma</i>	1–10,13,15–18	C	I,II	M,R,P,S,H
Blue-naped Chlorophonia	<i>Chlorophonia cyanea longipennis</i>	2,3,6,7,18	U	I	M,R,P,S,H
Chestnut-breasted Chlorophonia	<i>Chlorophonia pyrrhophrys</i>	8,9,15	SC	I	S

¹Letters in parenthesis refer to the global IUCN Red List categories: EN = Endangered, VU = Vulnerable, NT = Near threatened.

²Subspecies names are given, when available, from identified specimens in the museum or from unambiguous identifications from other evidence.

³Relative abundance: C = common (more than 10 individuals per day and daily encounters), FC = fairly common (less than 10 individuals per day and daily or almost daily encounters), U = uncommon (found in small numbers and not daily in prime habitat), SC = scarce (only occasionally found in small numbers), X = single record.

⁴Habitat: I = forest interior, II = edges and gaps, III = young second growth, pastures, and cultivated areas, IV = riverine vegetation or water courses, V = aerial or soaring.

⁵Type of evidence: M = museum specimen, R = tape recording, P = photograph, S = sight record, H = vocalization heard and confidently identified.

⁶Nearctic migrant.